





















Steller Sea Lion (SSLI)	Northern Fur Seal (NOFS)	Guadalupe Fur Seal (GUFS)
 <p>Adult female      Adult male</p>	 <p>Immature</p>	 <p>Immature</p>
<p><b>General Description:</b> body robust with long neck; ear pinna present; facial bones fuller and rounder; gap between upper 4<sup>th</sup> and 5<sup>th</sup> post-canines; color light tan to reddish brown; vibrissae smooth and beige (adult) or brown/beige (juvenile); adult males have full, furry neck/chest and a small sagittal crest. Very uncommon on mainland.</p>	<p><b>General Description:</b> body form long and slender with a long neck; ear pinna present (longer than sea lions); brown fur; buffy underparts; long guard hairs with very dense light-colored underfur (similar to sea otter); fur is soft and pliable and can lay in any direction; snout short, down-curved and pointed; head is football-shaped; vibrissae smooth and beige-colored; juveniles uncommon and adults very rare on mainland.</p>	<p><b>General Description:</b> body long and slender with long neck; ear pinna present (longer than sea lions); dark brown fur, buffy underparts; long guard hairs and very dense light-colored underfur (similar to sea otter); fur is soft and pliable and can lay in any direction; snout long and pointed with up-turned nose; juveniles uncommon and adults very rare on mainland. Does not breed in region, rare throughout CA.</p>
		
<p><b>Foreflipper:</b> Long and broad (compared to CSLI and fur seals) extending to the end of the ribs; fur on top of flipper; bottom bare; no nails.</p>	<p><b>Foreflipper:</b> long and narrow, extending to end of ribs; fur on top of front flipper stops at wrist (see arrow); top and bottom bare; no nails.</p>	<p><b>Foreflipper:</b> long and narrow, extending to the end of ribs; fur extending onto top surface of flipper (see arrow); bottom bare; no nails.</p>
		
<p><b>Hindflipper:</b> nails on digits 2-4; nails set back from tips by 1/4 the total length of flipper; fur on top of flipper, bottom bare.</p>	<p><b>Hindflipper:</b> extremely long and narrow; nails on digits 2-4; nails set back from tips by 1/5 the total length of flipper; “finger” tips of flipper extremely long (longer than both GUFS and CSLI); top of flipper furred, bottom bare.</p>	<p><b>Hindflipper:</b> long and narrow; nails on digits 2-4; nails set back from tips by 1/5 the total length of flipper; “finger” tips of flipper long (similar to CSLI but shorter than NOFS); top of flipper furred, bottom bare.</p>
<p><b>Size, Age, Sex:</b> sexual dimorphism (males larger); 100 cm (neonate/pup); 100 – 200cm (immature); 220 – 290 cm (adult female and sub-adult male); 240 – 330 cm (adult male).</p>	<p><b>Size, Age, Sex:</b> sexual dimorphism (males larger); 60 cm (neonate/pup); 100 cm (immature); 100 – 165 cm (adult female and sub-adult male); 190 – 230 cm (adult male).</p>	<p><b>Size, Age, Sex:</b> sexual dimorphism (males larger); 60 cm (neonate/pup); 100 – 150 cm (immature); 150 – 170 cm (adult female or sub-adult male); 180 – 240 cm (adult male).</p>

California Sea Lion (CSLI)	Harbor Seal (HASE)	Northern Elephant Seal (NESE)
 <p>Immature</p>  <p>Adult male</p>	 <p>Adult</p>	 <p>Immature</p>  <p>Adult male</p>
<p><b>General Description:</b> body form slender with long neck; ear pinna present; fur course and stiff with growth towards tail; color varies from honey to dark brown (based on age, sex, molt, and individual variation); vibrissae smooth and white (adult) or black (juvenile); yearling to adult common year-round.</p>	<p><b>General Description:</b> body rotund; short thick neck; pinna absent; snout blunt; color variable (white, silver or black with contrasting spots); fur coarse and stiff with growth towards the tail; vibrissae white and beaded; pups (Feb – April) and all age classes occur year-round in and around rookeries.</p>	<p><b>General Description:</b> body large and rotund; short thick neck; pinna absent; color gray to brown (black in pups) with no markings; fur coarse and stiff with growth towards the tail; vibrissae black and beaded. Adult males possess prominent proboscis and cracked, scarred chest. Common in or near rookeries from December - March. Molting season late Spring – Fall.</p>
 <p><b>Foreflipper:</b> long, extending to the end of the ribcage; fur on top surface of flipper; no nails; bottom surface bare.</p>	 <p><b>Foreflipper:</b> short and paddle-like, extending a few ribs past pit, back and front surfaces fully furred; nails present; digits similar in length.</p>	 <p><b>Foreflipper:</b> both sides of flipper fully furred, extending only a few ribs past pit; nails present; digit 1 is longer than others.</p>
 <p><b>Hindflipper:</b> nails on digits 2-4; nails set back from tips by 1/5 the total length of flipper; “finger” tips of flipper long (longer than GUFFS but not longer than NOFS); bottom surface bare.</p>	 <p><b>Hindflipper:</b> nails present on all digits; both front and back of flipper furred and webbed; digits 1-5 rounded.</p>	 <p><b>Hindflipper:</b> no nails on digits, both sides of flipper fully furred and webbed; digits 1 and 5 are much larger, longer and angled toward digits 2-4.</p>
<p><b>Size, Age, Sex:</b> Sexual dimorphism (males larger at maturity). 70 –90 cm (neonate/pup); 95 – 150 cm (immature); 150 – 180 cm (adult female or sub-adult male) 180 cm - 260 (adult male). Sexually mature males with prominent sagittal crest on top of skull; neck muscular and thick.</p>	<p><b>Size, Age, Sex:</b> No sexual dimorphism. 70 – 90 cm (pup); 90 – 150 cm (immature); 150 – 180 cm (adult).</p>	<p><b>Size, Age, Sex:</b> Extreme sexual dimorphism at maturity (males much larger); 120 – 140 cm (neonate “blackcoats”/pups 4-6 wks); 140 – 200 cm (immature); 200 – 320 cm (sub-adult male or adult female); 320 – 440 cm (adult male).</p>